


## Throwing Injuries in the Elbow

With the start of the baseball season each spring, doctors frequently see an increase in elbow problems in young baseball players. A common elbow problem in these children is medial apophysitis, commonly referred to by doctors as "Little Leaguer's elbow."

The elbow is the joint where the upper arm bone (humerus) meets the two bones of the lower arm (ulna and radius). The elbow is a combination hinge and pivot joint. The hinge part of the joint lets the arm bend like the hinge of a door; the pivot part lets the lower arm twist and rotate. Several muscles, nerves, and tendons (connective tissues between muscles and bones) cross at the elbow.



X-rays of the right elbow joint of an 8-year-old baseball pitcher show irregularity of the capitellum (humerus).

Injury occurs when repetitive throwing creates an excessively strong pull on the tendons and ligaments of the elbow. The young player feels pain at the knobby bump on the inside of the elbow.

"Little Leaguer's elbow" can be serious if it becomes aggravated. Repeated pulling can tear ligaments and tendons away from the bone. The tearing may pull tiny bone fragments with it in the same way a plant takes soil with it when it is uprooted. This can disrupt normal bone growth, resulting in deformity.

A less common condition called osteochondritis dissecans is also caused by excessive throwing, and may be the source of the pain on the outside of the elbow. Muscles work in pairs. In the elbow, if there is pulling on one side, there is pushing on the other side. As the elbow is compressed, the joint smashes immature bones together. This can loosen or fragment the bone and cartilage. The resulting condition is called osteochondritis dissecans.

### Risk Factors

"Little Leaguer's elbow" and osteochondritis dissecans affect pitchers and other players who throw repetitively. Continuing to throw may lead to major complications and jeopardize a child's ability to remain active in a sport that requires throwing.

The general guideline for how many pitches a child can safely throw is 200 in one week. This includes both practice and competitive play. To prevent throwing injuries, pitchers should only play three to four innings each game.

## Symptoms

"Little Leaguer's elbow" may cause pain on the inside of the elbow. A child should stop throwing if any of the following symptoms appear:

- Elbow pain
- Restricted range of motion
- Locking of the elbow joint

## Treatment Options

Left untreated, throwing injuries in the elbow can become a complicated condition. Younger children tend to respond better to nonsurgical treatments.

- Rest the affected area.
- Apply ice packs to bring down any swelling.
- If pain persists after a few days of complete rest of the affected area, or if pain recurs when throwing is resumed, stop the activity again until the child gets treatment.
- Refine throwing technique.

Surgery is occasionally necessary, especially in girls older than 12 years and boys older than 14 years.

Last reviewed and updated: July 2008



Reviewed by members of the Pediatric Orthopaedic Society of North America

*AAOS does not review or endorse accuracy or effectiveness of materials, treatments or physicians.*